



TECHNICAL NOTES



Popantel® F

Allwormer Tablets for Dogs / Large Dogs



Active Constituents

FEBANTEL, PYRANTEL and PRAZIQUANTEL

(10 kg dose: 250 mg, 49.8 mg and 50 mg per tablet respectively; 35 kg dose: 875 mg, 174.4 mg and 175 mg per tablet respectively)

Actions

Popantel® F is a treatment for gastrointestinal parasites of dogs. Popantel® F is effective against susceptible intestinal helminths. Popantel® F is not a preventative treatment for canine heartworm disease.

Indications

For the control in dogs of:

Roundworms (*Toxocara canis*, *Toxascaris leonina*)

Whipworms (*Trichuris vulpis*)

Hookworms (*Ancylostoma caninum*, *Uncinaria stenocephala*)

Tapeworms (*Dipylidium caninum* (common flea tapeworm), *Taenia ovis* (sheep measles), *T. pisiformis*, *T. hydatigena* and *Echinococcus granulosus* (hydatid tapeworm)).

Does not control heartworm.

Dosage and Administration

Popantel® F is given as a single dose by mouth immediately before or with a meal. Fasting prior to dosing is NOT necessary. IT IS ESSENTIAL TO WEIGH DOGS BEFORE TREATMENT. If worm problems persist, consult a veterinarian.

Jurox New Zealand Limited
8 Kordel Place, East Tamaki, Auckland, 2013 • Telephone: 0800 587 696 • Fax: 09 2948 012 • www.jurox.com
® Registered Trademark of Jurox Pty Limited

Products that work. People who care.





TECHNICAL NOTES



DOSAGE:

10 kg Popantel® F tablets

Bodyweight	Tablets
Up to 5 kg	½
5.1 – 10 kg	1
10.1 – 20 kg	2
20.1 – 30 kg	3
30.1 – 40 kg	4

35 kg Popantel® F tablets

Bodyweight	Tablets
20 - 35 kg	1
35.1 - 52 kg	1 ½
52.1 - 70 kg	2
70.1 - 87 kg	2 ½
Over 87 kg	3

PUPPIES: Treat at 2, 3, 4, 6, 9 and 12 weeks of age, then every 3 months.

ADULTS (over 6 months): Treat every 3 months. Dogs heavily infested with roundworms, hookworms or whipworms may require retreatment after 14 days. Treat every 4 weeks in sheep measles (*T. ovis*) infected locations.

PREGNANT / LACTATING BITCHES: Treat prior to mating, then 10 days before whelping and at 2 and 4 weeks after whelping.

First Aid

If poisoning occurs, contact a doctor or National Poisons Centre (0800 764 766).

Warning

Suspected of damaging fertility or the unborn child. May cause damage to the reproductive and the haemopoietic systems throughout prolonged or repeated exposure.

Handling precautions: Read label and obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Use personal protective equipment as required. Do not breathe dust. If medical advice is needed, have product container or label at hand.

Compatibility

Popantel® F may be combined with other treatments, for example heartworm prevention, flea or tick treatment.

Safety

Popantel® F may be given safely to all dogs including old animals, young pups, working dogs and greyhounds. It may be given to pregnant bitches but is only clinically indicated in the last 10 days prior to whelping.

Side effects: Vomiting may be observed in a small number of dogs after administration.



TECHNICAL NOTES



Disposal

Dispose of empty packaging in the household garbage.

Storage

Store locked up below 30°C (room temperature).

GHS Information

For GHS Information see Safety Data Sheet.

Presentation

For 10 kg Tablets: 8x10 tablets in cartons.

For 35 kg Tablets: 7x6 tablets in cartons.

Classification

Non-restricted Veterinary Medicine.

For animal treatment only.

Registration Numbers

ACVM Registration Nos. A010097 (dogs); A010098 (large dogs)